

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/10157 A1

- (51) International Patent Classification⁷: **C07D 405/06**, 405/14, A61K 31/505, A61P 31/04
- (21) International Application Number: **PCT/EP01/08426**
- (22) International Filing Date: **20 July 2001 (20.07.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
PCT/EP00/07357 **29 July 2000 (29.07.2000)** **EP**
- (71) Applicant (*for all designated States except US*): **ARPIDA AG** [CH/CH]; Dammstrasse 36, CH-4142 Münchenstein (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BURRI, Kaspar** [CH/CH]; Höhenweg 47, CH-4102 Binningen (CH). **GILLESSEN, Dieter** [CH/CH]; Oberfeldstrasse 12, CH-4133 Pratteln (CH). **GREIVELDINGER-POE-NARU, Sorana** [FR/CH]; Thermenstrasse 19, CH-4310 Rheinfelden (CH). **ISLAM, Khalid** [GB/CH]; Binningerstrasse 82, CH-4153 Reinach (CH).
- (74) Agent: **HOFMANN, Dieter**; StratAll, Therwilerstrasse 87, CH-4153 Reinach (CH).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/10157 A1

(54) Title: **BENZOFURAN DERIVATIVES AND THEIR USE AS ANTIBACTERIAL AGENTS**

(57) Abstract: The invention relates to novel benzofuran derivatives and their use as active ingredients in the preparation of pharmaceutical compositions. The invention also concerns related aspects including processes for the preparation of the compounds, pharmaceutical compositions containing one or more of those compounds and especially their use as anti-infectives.

BENZOFURAN DERIVATIVES AND THEIR USE AS ANTIBACTERIAL AGENTS

10 The present invention relates to novel 2,4-diamino-5- (substituted) pyrimidines, to pharmaceutical compositions containing them, to processes for preparing them and their compositions, to intermediates for making them and to their use in the treatment of microbial infections.

15 Certain 2,4-diamino-5-benzylpyrimidines have been demonstrated to be potent inhibitors of dihydrofolate reductase (DHFR), which catalyses the reduction of dihydrofolic acid to tetrahydrofolic acid (THFA). This property has been shown to result frequently in useful pharmaceutical properties particularly in the treatment of bacterial infections. Thus, U.K. Patent Specification No. 875,562 discloses *inter alia* 2,4-diamino-5-benzylpyrimidines wherein the benzyl moiety is substituted by three
20 C₁₋₄ alkoxy groups.

Trimethoprim, 2,4-diamino-5-(3,4,5-trimethoxybenzyl)pyrimidine, is specifically disclosed in U.K. Patent No. 875, 562 and is the most active general antibacterial agent amongst the 2,4-diamino-5-benzylpyrimidines known to date. Due to their mode
25 of action, these benzylpyrimidines potentiate the antibacterial activity of the sulphonamides, and Trimethoprim has been used extensively over the last decade in human therapy in combination with various sulphonamides, and in particular with sulphamethoxazole, for the treatment of bacterial infections.

European Patent Applications Nos. 81109631.2 and 83104240.3 disclose *inter alia*
30 also such type of compounds and their use.


It has now been found that a group of novel benzofuran derivatives is more potent than e. g. Trimethoprim and is especially active against Gram positive pathogens like *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Enterococcus faecalis* or
35 *Streptococcus pneumoniae* and at the same time also against Gram negative pathogens like *Haemophilus influenzae*, *Escherichia coli*, *Klebsiella pneumoniae* or

Proteus vulgaris. The compounds proved to be especially potent against respiratory tract pathogens.

Therefore, the present invention relates to novel compounds of the general formula I

5

Formula I



wherein

- 10 R¹ represents straight or branched chain lower alkyl with 2 to 6 carbon atoms; cycloalkylmethyl with 3 to 6 carbon atoms; aryl; arylmethyl or heteroarylmethyl, the aryl and heteroaryl group may be mono-, di- or tri- substituted with halogen, amino, hydroxy, nitro, trifluoromethyl, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; straight
- 15 or branched chain lower alkylcarbonyl with up to 6 carbon atoms; cycloalkylcarbonyl with 3 to 6 carbon atoms; cycloalkylhydroxymethyl with 3 to 6 carbon atoms; arylcarbonyl, the aryl group may be mono-, di or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; arylhydroxymethyl, the aryl group may be
- 20 mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; straight or branched chain lower alkenyl with 2 to 6 carbon atoms; hydroxy-lower alkyl with 1 to 6 carbon atoms; fluoro-lower alkyl with 1 to 6 carbon atoms; aryloxy-lower alkyl whereby the aryl group may be mono-, di- or tri- substituted
- 25 with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; arylthio-lower alkyl whereby the aryl group may be mono-, di- or tri- substituted with halogen, amino, lower

alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; arylamino-lower alkyl whereby the aryl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy; lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; lower alkenyloxy lower alkyl whereby the lower alkenyl group may contain 2 to 4 carbon atoms and the lower alkyl group may contain 1 or 2 carbon atoms; benzyloxy lower alkyl whereby the benzyl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; lower alkylamino lower alkyl whereby the lower alkyl groups may contain 1 to 3 carbon atoms; heterocyclylmethyl containing one to three hetero atoms which can be the same or different and which may be substituted with lower alkyl, halogen, amino, lower alkyloxy, hydroxy, lower alkylcarbonylamino, arylcarbonylamino and benzofused derivatives thereof.

R^2 and R^3 independently represent hydrogen; lower alkyl with 1 to 3 carbon atoms; or together a lower alkylene group with 1 to 3 carbon atoms bridging the oxygen atoms and forming a five, six or seven membered ring;

R^4 represents hydrogen; straight or branched chain lower alkyl with 1 to 4 carbon atoms;

and pharmaceutically acceptable salts and N-oxides thereof.

In the definitions of the general formula I – if not otherwise stated – the expression lower means straight and branched chain groups with either one or two to six or three carbon atoms, preferably 1 to 3 carbon atoms. Examples of lower alkyl and lower alkoxy groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec.-butyl, tert.-butyl, pentyl, hexyl, trifluoromethyl, chloromethyl, fluoromethyl, hydroxymethyl, carbahdehyd, thiomethyl, methoxy, ethoxy, propoxy, butoxy, iso-butoxy, sec.-butoxy and tert.-butoxy. Lower alkylene groups as bridging two oxygen atoms are preferably methylen(-dioxy), ethylen(-dioxy) and propylen(-dioxy) groups and forming in this way a five-, six- or seven-membered ring. Examples of lower alkanoyl-groups are acetyl, propanoyl and butanoyl. Lower alkenylen means e.g. vinylen, propenylen and butenylen. Lower alkenyl and lower alkynyl means groups like ethylen, propylen, butylen, 2-methyl-propenyl, and ethinylen, propinylen, butinylen, pentinylen, 2-methyl-pentylen etc. Lower alkenyloxy means allyloxy, vinyloxy, propenyloxy and the like.

The expression **cycloalkyl** means a saturated cyclic hydrocarbon ring with 3 to 6 carbon atoms, e.g. cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, which may be substituted with lower alkyl, hydroxy-lower alkyl, amino-lower alkyl, lower alkoxy-lower alkyl and lower alkenylen groups. The expression **heteroaryl** means six-membered aromatic rings containing one to four nitrogen atoms, benzofused six-membered aromatic rings containing one to three nitrogen atoms, five-membered aromatic rings containing one oxygen or one nitrogen or one sulfur atom, benzo-fused five-membered aromatic rings containing one oxygen or one nitrogen or one sulfur atom, five membered aromatic rings containig an oxygen and nitrogen atom and benzo fused derivatives thereof, five-membered aromatic rings containing a sulfur and a nitrogen atom and benzo fused derivatives thereof, five-membered aromatic rings containing two nitrogen atoms and benzo fused derivatives thereof, five membered aromatic rings containing three nitrogen atoms and benzo fused derivatives thereof or the tetrazolyl ring e.g. furanyl, thienyl, pyrrolyl, pyridinyl, indolyl, quinolinyl, isoquinolinyl, imidazolyl, triazinyl, thiazinyl, thiazolyl, isothiazolyl, pyridazinyl, oxazolyl, isoxazolyl, etc. whereby such rings may be substituted with lower alkyl, lower alkenyl, amino, amino-lower alkyl, halogen, hydroxy, lower alkoxy, trifluoromethoxy or trifluoromethyl. The expression **aryl** represents unsubstituted as well as mono-, di- or tri-substituted aromatic rings with 6 to 10 carbon atoms like phenyl or naphthyl rings which may be substituted with aryl, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, lower alkenyloxy, lower alkynyl-lower alkyl-oxy, lower alkenylen, lower alkyleneoxy or lower alkylendioxy forming with the phenyl ring a five- or six-membered ring, hydroxy-lower alkyl, hydroxy-lower alkenyl, hydroxy-lower alkyl-lower alkynyl, lower alkyloxy-lower alkyl, lower alkyloxy-lower alkyloxy, trifluoromethyl, trifluoromethoxy, cycloalkyl, hydroxy-cycloalkyl, heterocyclyl, heteroaryl. The expression **heterocyclyl** represents saturated and unsaturated, but not aromatic, three- to six-membered rings containing one to three nitrogen, oxygen or sulfur atoms which may be the same or different like azyridinyl, piperidinyl, mopholinyl, piperazinyl, tetrahydropyranyl, dihydropyranyl, 1,4-dioxanyl, pyrrolidinyl, tetrahydrofuranyl, dihydroimidazolyl, dihydropyrazoyl, pyrazolidinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, dihydroquiniliny, dihydroisoquinolinyl.

Preferred compounds are compounds of formula I, wherein R¹ is straight or branched chain lower alkyl with 2 to 6 carbon atoms; cycloalkylmethyl with 3 to 6 carbon atoms; aryl; arylmethyl or heteroarylmethyl, the aryl and heteroaryl group may be mono- or di substituted with halogen, amino, hydroxy, nitro, trifluoromethyl, lower alkyloxy, lower

alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different, and R^2 and R^3 are methyl or together are a methylene group bridging the oxygen atoms to which they are attached and R^4 is hydrogen or methyl.

5 Especially preferred compounds are compounds of formula I, wherein R^1 is 4-methoxy-benzyl, phenyl, benzyl, cyclopropylmethyl, 4-fluoro-benzyl, 3,4-dihydro-1H-isoquinolin-2-ylmethyl, 4-methoxy-benzylloxymethyl, 4-acetylamino-phenyl-sulfanyl-methyl, 4-trifluoromethyl-phenoxy-methyl, 4-amino-phenoxy-methyl, allyloxymethyl, phenyl-[1,3]dioxolan-2-yl, pyrrol-1-ylmethyl, 3,4-dimethoxy-benzyl, 4-
10 hydroxyphenylloxymethyl, phenylcarbonylmethyl, 4-fluoro-phenoxy-methyl, 2,2-dimethyl-propyl, 4-fluoro-phenylsulfanylmethyl, hydroxymethyl, formyl, 4-fluoro-phenylamino-methyl, imidazol-1-ylmethyl, dimethylaminomethyl, morpholin-4-ylmethyl, biphenyl-4-ylmethyl, ethoxycarbonyl, carboxy, 4-hydroxybenzyl, furan-2-ylmethyl, thiophen-2-ylmethyl, thiophen-3-ylmethyl, 1-methyl-pyrrol-2-ylmethyl, pyridin-2-
15 ylmethyl, pyridin-3-ylmethyl, pyridin-4-ylmethyl, 4-acetylamino-benzyl, 4-amino-benzyl, 4-nitro-benzyl, 4-acetylamino-3-methoxy-benzyl, 4-amino-3-methoxy-benzyl, 3-acetylamino-4-methoxy-benzyl, 3-amino-4-methoxy-benzyl, allyl, isopropenyl or halogenmethyl, and R^2 and R^3 are methyl or together are a methylene group bridging the oxygen atoms to which they are attached and R^4 is hydrogen or methyl.

20

Very preferred compounds are compounds of formula I, wherein R^1 is 4-methoxy-benzyl, phenyl, benzyl, cyclopropylmethyl, 4-fluoro-benzyl, 3,4-dihydro-1H-isoquinolin-2-ylmethyl, 4-methoxy-benzylloxymethyl, 4-acetylamino-phenyl-sulfanyl-methyl, 4-trifluoromethyl-phenoxy-methyl, 4-amino-phenoxy-methyl, allyloxymethyl, 3,4-
25 dimethoxy-benzyl, 4-hydroxybenzyl, pyridin-2-ylmethyl, pyridin-3-ylmethyl, pyridin-4-ylmethyl, 4-acetylamino-benzyl, 4-amino-benzyl, 4-acetylamino-3-methoxy-benzyl, 4-amino-3-methoxy-benzyl, 3-acetylamino-4-methoxy-benzyl, 3-amino-4-methoxy-benzyl or aryl methyl and R^2 and R^3 are both methyl and R^4 is hydrogen.

30 Most preferred compounds are compounds of formula I, wherein R^1 is 4-methoxy-benzyl, benzyl, cyclopropylmethyl, 4-fluoro-benzyl, 3,4-dihydro-1H-isoquinolin-2-ylmethyl, 4-amino-phenoxy-methyl, 4-hydroxybenzyl, pyridin-2-ylmethyl, pyridin-3-ylmethyl, pyridin-4-ylmethyl, 4-acetylamino-benzyl, 4-amino-benzyl, 4-acetylamino-3-methoxy-benzyl, 4-amino-3-methoxy-benzyl, 3-acetylamino-4-methoxy-benzyl, 3-
35 amino-4-methoxy-benzyl and R^2 and R^3 are both methyl and R^4 is hydrogen

Preferred compounds of the present invention include:

- 5-[6,7-Dimethoxy-2-(4-methoxy-benzyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5 5-(6,7-Dimethoxy-2-phenyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(2-Benzyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(2-Cyclopropylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-
- 10 diamine,
- 5-[2-(4-Fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-[2-(3,4-Dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-
- 15 pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 20 N-{4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethylsulfanyl]-phenyl}-acetamide,
- 5-[6,7-Dimethoxy-2-(4-trifluoromethyl-phenoxy-methyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 25 5-[2-(4-Amino-phenoxy-methyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-(2-Allyloxymethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 30 5-[6,7-Dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-(6,7-Dimethoxy-2-pyrrol-1-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 35

- 5-[2-(3,4-Dimethoxy-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethoxy]-phenol,
- [4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-phenyl-methanone,
- 5-[2-(4-Fluoro-phenoxy-methyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-[2-(2,2-Dimethyl-propyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-[2-(4-Fluoro-phenylsulfanylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- [4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-methanol,
- 4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-carbaldehyde
- 5-{2-[(4-Fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-ylmethyl}-pyrimidine-2,4-diamine,
- 5-(2-Imidazol-1-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(2-Dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(6,7-Dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(2-Biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,

4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid,

5 4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid ethyl ester,

5-[6,7-Dimethoxy-2-((4-chlorobenzyl)-benzofuran-4-ylmethyl)]pyrimidine-2,4-diamine,

10 5-[6,7-Dimethoxy-2-(1-naphthylmethyl)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(2-propenyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

15 5-(6,7-Dimethoxy-2-trifluoromethylbenzofuran-4-ylmethyl)pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(2,2-dimethylpropan-1-one)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

20 5-[6,7-Dimethoxy-2-(cyclopropylcarbonyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(4-methoxyphenyl-methanone)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

25 5-[6,7-Dimethoxy-2-(4-chlorophenyl-methanone)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(4-fluorophenyl-methanone)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

30 5-[6,7-Dimethoxy-2-(1-naphthyl methanone)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

35 5-[6,7-Dimethoxy-2-(2,2-dimethyl-1-hydroxypropyl)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(cyclopropylmethanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(phenylmethanol)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5

5-[6,7-Dimethoxy-2-((4-methoxyphenyl)methanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-((4-chlorophenyl)methanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

10

5-[6,7-Dimethoxy-2-((4-fluorophenyl)methanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

15 5-[6,7-Dimethoxy-2-(1-naphthylmethanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[2-(3,4-Dihydro-2H-quinolin-1-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,

20

5-[6,7-Dimethoxy-2-(tetrazol-5-ylmethyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(indol-1-ylmethyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

25

5-[6,7-Dimethoxy-2-(imidazol-1-ylcarbonyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-(2-Furan-2-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,

30

5-(6,7-Dimethoxy-2-thiophen-2-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,

5-[6,7-Dimethoxy-2-(1-methyl-1H-pyrrol-2-ylmethyl)-benzofuran-4-ylmethyl]-

35

pyrimidine-2,4-diamine,

- 5-(6,7-Dimethoxy-2-pyridin-2-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
5-(6,7-Dimethoxy-2-pyridin-3-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
5 5-(6,7-Dimethoxy-2-pyridin-4-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-phenol,
N-{4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-
10 phenyl}-acetamide,
5-[2-(4-Amino-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
5-[6,7-Dimethoxy-2-(4-nitro-benzyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
15 N-{4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-2-
methoxy-phenyl}-acetamide,
5-[2-(4-Amino-3-methoxy-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-
20 2,4-diamine,
N-{5-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-2-
methoxy-phenyl}-acetamide,
25 5-[2-(3-Amino-4-methoxy-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-
2,4-diamine,
5-(6,7-Dimethoxy-2-thiophen-3-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-
30 diamine,
and pharmaceutically acceptable salts and N-oxides thereof.

The expression pharmaceutically acceptable salts encompasses either salts with inorganic acids or organic acids like hydrohalogenic acids, e.g. hydrochloric or
35 hydrobromic acid; sulfuric acid, phosphoric acid, nitric acid, citric acid, formic acid, acetic acid, maleic acid, tartaric acid, methanesulfonic acid, p-toluenesulfonic acid

and the like or in case the compound of formula I is acidic in nature with an inorganic base like an alkali or earth alkali base, e.g. sodium hydroxide, potassium hydroxide, calcium hydroxide etc.

Because of their ability to inhibit Gram positive and Gram negative bacteria, the
5 described compounds can be used for the treatment of diseases which are associated with an infection by such type of pathogens. They are valuable anti-infectives.

The compounds can be administered orally, rectally, parenterally, e.g. by intravenous,
10 intramuscular, subcutaneous, intrathecal or transdermal administration or sublingually or as ophthalmic preparation or administered as aerosol. Examples of applications are capsules, tablets, orally administered suspensions or solutions, suppositories, injections, eye-drops, ointments or aerosols/nebulizers.

15 Preferred applications are intravenous, intra-muscular, or oral administrations as well as eye drops. The dosage used depends upon the type of the specific active ingredient, the age and the requirements of the patient and the kind of application. Generally, dosages of 0.1 – 50 mg / kg body weight per day are considered. The preparations with compounds of formula I can contain inert or as well
20 pharmacodynamically active excipients like sulphonamides. Tablets or granules, for example, could contain a number of binding agents, filling excipients, carrier substances or diluents.

These compositions may be administered in enteral or oral form e.g. as tablets,
25 dragees, gelatine capsules, emulsions, solutions or suspensions, in nasal form like sprays or rectally in form of suppositories. These compounds may also be administered in intramuscular, parenteral or intravenous form, e.g. in form of injectable solutions.

30 These pharmaceutical compositions may contain the compounds of formula I as well as their pharmaceutically acceptable salts in combination with inorganic and/or organic excipients which are usual in the pharmaceutical industry like lactose, maize or derivatives thereof, talcum, stearic acid or salts of these materials.

For gelatine capsules vegetable oils, waxes, fats, liquid or half-liquid polyols etc. may be used. For the preparation of solutions and syrups e.g. water, polyols, saccharose, glucose etc. are used. Injectables are prepared by using e.g. water, polyols, alcohols, glycerin, vegetable oils, lecithin, liposomes etc. Suppositories are prepared by using
5 natural or hydrogenated oils, waxes, fatty acids (fats), liquid or half-liquid polyols etc.


The compositions may contain in addition preservatives, stabilisation improving substances, viscosity improving or regulating substances, solubility improving substances, sweeteners, dyes, taste improving compounds, salts to change the
10 osmotic pressure, buffer, antioxidants etc.

The compounds of formula I may also be used in co-therapy with one or more other therapeutically used classes of antimicrobial substances, for example, β -lactams e.g. penicillins and cephalosporins; glycopeptides; quinolones; tetracyclines;
15 aminoglycosides; macrolides etc.

The dosage may vary within wide limits but should be adapted to the specific situation. In general the dosage given in oral form should daily be between about 3 mg and about 4 g, preferably between about 0.2 g and about 4 g, especially preferred
20 between 0.2 g and 2 g per adult with a body weight of about 70 kg. The dosage should be administered preferably in 1 to 3 doses per day which are of equal weight. As usual children should receive lower doses which are adapted to body weight and age.

The invention also relates to a process for the manufacture of compounds of formula I

Formula I



I


5

wherein

R^1 , R^2 , R^3 and R^4 have the meaning given in formula I above,

which process comprises reacting – as depicted in **Scheme 1** – a compound of the
 10 general formula III (obtained from a gallate derivative II: either by Friedel-Crafts
 acylation, or via a Vilsmeier aldehyde synthesis (K. Hayashi, K. Tokura, K. Okabe, K.
 Yamamoto, and K Tawara, *Chem. Pharm. Bull.* **1982**, *30*, 2860-2869), or by
 formylation with dichloromethoxymethane, see experimental part), with boron
 trichloride or boron tribromide at temperatures between -70°C and 0°C . The methoxy
 15 group adjacent to the acyl group is thereby selectively cleaved, to yield ortho-acyl
 phenol derivatives of the general formula IV:

Scheme 1



20


Subsequently, the compounds **IV**, wherein R^2 , R^3 , and R^4 have the meaning given in formula I above, are reacted with α -bromomethyl ketones R^5 -CO-CH₂Br **V** in the presence of potassium carbonate, to yield 2-acylated benzofuran derivatives **VI** (**Scheme 2**)

5

whereby in formula **V** R^5 represents straight or branched chain lower alkyl with 1 to 5 carbon atoms; cycloalkyl with 3 to 6 carbon atoms; aryl or heteroaryl, the aryl and heteroaryl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents
10 may be the same or different; straight or branched chain lower alkenyl with 2 to 6 carbon atoms.

The carbonyl group of **VI** can then be removed selectively by reducing it with sodium cyanoborohydride in the presence of trimethylsilyl chloride, to yield the benzofuran
15 ester derivatives **IX**, wherein R^1 , R^2 , R^3 , and R^4 have the meaning given in formula I.


Scheme 2



Alternatively, benzofurans of the general structure **IX** can be obtained by reacting **IV** with an α -bromo-ester derivative $R^6\text{OCO-CH(Br)-R}^7$ **VII** (R^6 is *tert*-butyl, and R^7 may be hydrogen or *tert*-butoxycarbonyl), again in the presence of carbonate, to give benzofuran dicarboxylates of the formula **VIII**, whereby R^2 , R^3 , and R^4 have the meaning given in formula **I**, and R^6 represents the *tert*-butyl group. The latter can be cleaved selectively with trifluoroacetic acid, and may then be reduced preferentially with diborane. The resulting primary alcohol is then converted into a leaving group, e.g. a tosylate, which is then displaced by a nucleophile R^8 , such as a nitrogen-containing heteroaromatic moiety to yield **IX** wherein R^1 equals $R^8\text{CH}_2$ -.


If the acyl group in formula **VI** (**Scheme 2**) is to be preserved, i.e. if in the final product (formula **I**) R^1 represents an acyl group (cycloalkylcarbonyl, arylcarbonyl etc.), it may be protected as an ethylene ketal, as set forth in structure **X** (**Scheme 3**):

15


Scheme 3

In a third alternative approach described in **Scheme 4**, phenol **XI** (Wipf, P.; Weiner, W.S. *J. Org. Chem.* **1999**, *64*, 5321-5324) is reacted with a propargylic alcohol derivative **XII** under Mitsunobu conditions (diethyl azodicarboxylate and triphenyl phosphine) to afford the phenyl ethers **XIII**. After cleaving off the trimethylsilyl protecting group with potassium carbonate, the resulting free acetylenes can be rearranged, as described previously (Koch-Pomeranz, U.; Hansen, H.-J.; Schmid, H. *Helv. Chim. Acta* **1973**, *56*, 2981-3004), to benzofurans **IX**.

Scheme 4



Scheme 5




XIV = IX, X, XXVII or XIX

- 5 The manufacturing process then continues as depicted in **Scheme 5**. The ester function of the general substructure **XIV** (consisting of either **IX** or **X**) is reduced to an aldehyde **XVI**, either directly, with an organoaluminum hydride reagent, or in two steps, via the primary alcohol **XV**.


Finally, the aldehyde group is transformed into the diaminopyrimidine ring, employing the standard technology outlined in **Scheme 6**.

5

Scheme 6

Thus, compounds of formula I can be obtained; if R^1 represents an acyl group, i.e. structure **XVII** has been the intermediate, the final elaboration includes an acid-induced removal of the ketal protecting group.

15

Scheme 7


Access to an alternative array of substituents can be gained by proceeding according to **Scheme 7**: The diester **VIII** (**Schemes 2 and 7**), if R^6 represents a tert.-butyl group, can be cleaved selectively with trifluoroacetic acid, yielding **VIII**, $R^6=\text{H}$. This carboxylic acid may then be reduced with diborane to the primary alcohol **IX**, $R^1=\text{CH}_2\text{OH}$.

Treatment with thionyl chloride affords IX, $R^1=CH_2Cl$. This very active halide can now be employed as one component in the Lewis acid (preferably $ZnCl_2$) catalysed Friedel-Craft's alkylation, the R^5 component being either an electron-rich aryl or heteroaryl group which may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, such as phenol, 1,2-dimethoxybenzene, 2-methoxyphenacetin, furan, thiophene, 1-methylpyrrol. The product is compound IX, wherein R^1 , R^2 , R^3 , and R^4 have the meaning given in formula I

Yet another alternative to 2-substituted benzofurans is described in Scheme 8:

Here the same ester VIII, $R^6=tert.-butyl$, is cleaved with trifluoroacetic acid, and the resulting carboxylic acid VIII, $R^6=H$ can be coupled in peptide fashion with primary or secondary amines R^8 , producing the carboxamides XVIII, wherein R_8 may be a mono- or di-lower alkylamino group, aminoaryl or heterocycle acylated at the secondary amine group, such as morpholin-4-yl or tetrahydroisoquinoline-2-yl and R^2 , R^3 , and R^4 have the meaning given in formula I.

Scheme 8




A different pattern of substituents results, when the free acid **VIII**, $R^6=H$ is reduced, as in **Scheme 7**, to the primary alcohol **XIX**, $R^9=H$ (this compound is identical with **IX**, $R^1=CH_2OH$ of **Scheme 7**). This alcohol in turn may be transformed into ether derivatives **XIX** by alkylating its sodium salt with very reactive halides such as allyl bromide or benzyl bromide, wherein R^9 may be lower alkyl, lower alkenyl, or arylmethyl and R^2 , R^3 , and R^4 have the meaning given in formula **I**

The final elaboration of the intermediates **IX**, **XVIII** and **XIX** of the **Schemes 7** and **8** into diaminopyrimidines is then carried out in analogy to the **Schemes 5** and **6**.

If so desired, the operations summarised in the **Schemes 7** and **8** may be carried out in a more convergent fashion closer to the end of the synthesis:


15

Scheme 9

Thus, as in **Scheme 9**, from the allyl protected product **XX** (obtained from **XIX**, $R^9 =$ allyl, in **Scheme 8** by construction of the pyrimidine ring according to **Schemes 5** and **6**) the hydroxymethyl group can be liberated by isomerising of the allyl group with $Pd(PPh_3)_4$ and then hydrolizing the resulting enol ether. Treatment with thionyl chloride affords the reactive chloride **XXI**, which, in turn can be employed as a Friedel-Craft's component as described in **Scheme 7**, or which can be used to O- N- or S-alkylate with HXR^{10} , wherein X may be one oxygen or one nitrogen or one sulfur and R^{10} alkyl or aryl groups such as phenols, primary or secondary alkyl- or dialkylamines and aryl(alkyl)amines, thiophenols, pyrrol, indol, imidazol or tetrazol to yield compounds of formula I.

An even more convergent approach to is outlined in **Scheme 10**:


Scheme 10



Here, the demethylated phenol **XXII** (G. Rey-Bellet and R. Reiner *Helv. Chim. Acta* 1970, 53, 945) is treated with α -chlorinated aldehydes (or ketones) **XXIII** wherein R^1 and R^4 have the meaning given in formula I, in the presence of potassium carbonate, yielding the final products I directly.

Another convergent approach (**Scheme 11**) consists in brominating the phenol **XXIV** and coupling the resulting bromophenol **XXV** with acetylenic component **XXVI** as described in the literature (*J. Org. Chem.* 1996, 61, 9280-9288; *Bioorg. Med. Chem.* 1999, 7, 6, 1131-1140).

Scheme 11



5

The coupled acetylene then closes spontaneously to the corresponding benzofuran XXVII.

Examples

Abbreviations:

- ACN: Acetonitrile
- 5 ATCC: American type culture collection
- DEAD: Diethyl azodicarboxylate
- DMF: Dimethyl formamide
- DMSO: dimethyl sulfoxide
- EtOH: Ethanol
- 10 ESI: Electrospray ionisation
- FC: Flash chromatography
- HPLC: High performance liquid chromatography
- MeOH: methanol
- MS: Mass spectrometry
- 15 NMR: Nuclear magnetic resonance
- RedAl: Sodium bis(2-methoxyethoxy)aluminum hydride
- tBuOK: Potassium tert-butoxide
- TBME: tert-Butyl methyl ether
- TFA : Trifluoroacetic acid
- 20 THF: Tetrahydrofuran
- TLC: Thin layer chromatography
- TMSCl: trimethyl chlorosilane

General procedure A: Formylation

- 25 Under argon at -20°C , to a solution of compound II (1.0 eq.) in dry CH_2Cl_2 was added dichloromethylmethyl ether (1.5 eq.). SnCl_4 (1.0 eq.) was added dropwise over a period of 30 min, while maintaining the inside temperature at -20°C . After the addition, the mixture was allowed to slowly warm up to room. After disappearance of the starting material (TLC), the reaction mixture was again cooled to -10°C and
- 30 quenched by slow addition of NaHCO_3 . The organic phase was collected, the aqueous phase was extracted with three portions of tert-butyl methyl ether (TBME), the combined organic phases were washed with brine, dried with MgSO_4 and the solvents removed under reduced pressure to give compound III.

Example 1:

Methyl 2-formyl-3,4,5-trimethoxybenzoate was obtained as a yellow foam (39.0 g, 90%, 85% HPLC purity) by reacting methyl 3,4,5-trimethoxybenzoate (36.0 g) with dichloromethylmethyl ether (27.0 g, 1.5 eq.) and SnCl₄ (41.5 g, 1.0 eq). The product, which is rather unstable, was directly used for the next step.

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 10.28 (s, 1H, COH), 6.93 (s, 1H, Ar), 3.96 (s, 3H, OMe), 3.93 (s, 3H, OMe), 3.89 (s, 6H, 2 OMe).

10 General procedure B: Demethylation

Under argon at -10°C, to a solution of compound III (1 eq.) in methylene chloride, BBr₃ (0.5 eq.) was added over a period of 20 min, such that the temperature was not exceeding 0°C. The mixture was stirred at 0°C, until disappearance of the starting material. The reaction mixture was poured into ice-water and the aqueous layer was extracted with two portions of TBME. The combined organic phases were washed with water, dried with MgSO₄, and the solvents evaporated under reduced pressure to give compound IV.

Example 2:

20 Methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (36.8 g, 98%, 88% HPLC purity) was obtained as a white crystalline compound by reacting methyl 2-formyl-3,4,5-trimethoxybenzoate (39.5 g) with BBr₃ (7.5 ml, 0.5 eq.).

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 12.28 (s, 1H, OH), 10.26 (s, 1H, COH), 6.57 (s, 1H, Ar), 4.01 (s, 3H, OMe), 3.98 (s, 6H, 2 OMe).

25

General procedure C: Synthesis of benzofurans

Under argon, compound IV (1 eq.) was dissolved in DMF or toluene (Fluka on 4Å Mol. sieve), K₂CO₃ powder (2.4 eq.) and in some cases tetrabutylammonium bromide (0.1 eq.) was added. After 10 min stirring at r.t., a bromoketone V, a bromomalonic ester VII or a α -bromoester VII (1.2 to 1.9 eq.) was added as well as some 4Å Mol. sieves. The reaction was stirred 3h to 24h at 110°C. The reaction mixture was filtrated and the solvent evaporated. The residual brown oil was dissolved in CH₂Cl₂ and washed with water and brine. The organic layer was dried with MgSO₄, the solvent evaporated to give the compound VI or VIII.

Example 3:

6,7-Dimethoxy-2-cyclopropylcarbonyl-benzofuran-4-carboxylic acid methyl ester (128 mg, 59%) was obtained as a yellow solid after crystallisation from ethyl acetate by reacting methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (170 mg, 0.70 mmol) with
5 2-bromo-1-cyclopropylethanone (0.138 g, 0.849 mmol) (which was previously obtained from the reaction of 1-cyclopropyl-vinyltrimethylsilane with N-bromosuccinimide), K₂CO₃ (170 mg, 1.7 eq.) and tetrabutylammonium bromide (27 mg, 0.07 mmol)

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 7.98 (s, 1H, Ar), 7.68 (m, 1H, Ar), 4.28 (s, 3H, OMe), 3.93 (s, 3H, OMe), 3.92 (s, 3H, OMe), 2.72-2.55 (m, 1H), 1.29-1.25 (m, 2H),
10 1.09-1.04 (m, 2H).

Example 4:

6,7-Dimethoxy-2-phenylcarbonyl-benzofuran-4-carboxylic acid methyl ester (1.1 g, 56%) was obtained as a yellow solid by reacting Methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (1.5 g, 6.24 mmol) with K₂CO₃ powder (2.07 mg, 14.97 mmol) and phenacyl bromide (2.36 mg, 11.86 mmol)
15

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 8.04 (s, 1H, Ar), 7.74 (s, 1H, Ar), 7.68-7.54 (m, 5H, Ar), 4.37 (s, 3H, OMe), 4.00 (s, 3H, OMe), 3.97 (s, 3H, OMe).

20 MP: 116-120°C

Example 5:

6,7-Dimethoxy-2-(2,2-dimethylpropanoyl)-benzofuran-4-carboxylic acid methyl ester (182 mg, 57%) was obtained as a yellow solid by reacting methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (250 mg, 1.04 mmol) with K₂CO₃ powder (338 mg, 2.44 mmol) and 1-bromopinacol (265 μ l, 1.96 mmol).
25

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 8.04 (s, 1H, Ar), 7.72 (s, 1H, Ar), 4.35 (s, 3H, OMe), 3.98 (s, 3H, OMe), 3.96 (s, 3H, OMe), 1.45 (s, 9H, tert-Bu)

MP 104-108°C

30

Example 6:

6,7-Dimethoxy-2-tert-butoxycarbonyl-benzofuran-4-carboxylic acid methyl ester (100 mg, 37%) was obtained as a yellow solid by reacting methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (200 mg, 0.83 mmol) with K₂CO₃ powder (172 mg, 1.24 mmol),
35 tetrabutylammonium bromide (27 mg, 0.08 mmol) and bromomalonic acid tert-butyl ester (294 mg, 1.00 mmol).

General procedure D: Reduction of the ketone

Under argon at 0°C, to a solution of the ketone VI (1 eq.) in freshly distilled THF trimethylsilyl chloride (10 eq.), 4Å Mol sieve powder and then sodium cyanoborohydride (10 eq.) were added. The reaction was complete after stirring it for 5h to 24 h at r.t. Dichloromethane was added to the reaction mixture, which was then filtrated through celite. After several washings of the residue with CH₂Cl₂, the filtrate was washed with water and brine, dried with MgSO₄ and the solvents evaporated to give compound IX.

Example 7:

6,7-Dimethoxy-2-cyclopropylmethyl-benzofuran-4-carboxylic acid methyl ester (94 mg, 90%) was obtained as a glassy solid by reacting trimethylsilyl chloride (46 µl, 3.61 mmol) and sodium cyanoborohydride (240 mg, 3.61 mmol).

MS ESI: 291 (M+H)

15

Example 8:

6,7-Dimethoxy-2-phenylmethyl-benzofuran-4-carboxylic acid methyl ester (186 mg, 95%) was obtained as a glassy solid, after purification by flash chromatography (R_f 0.66, 3:2, hexane/ethyl acetate) by reacting 6,7-dimethoxy-2-phenylcarbonyl-benzofuran-4-carboxylic acid methyl ester (200 mg, 0.588 mmol) with sodium cyanoborohydride (370 mg, 5.88 mmol) and trimethylsilyl chloride (743 µl, 5.88 mmol).

MS ESI+: 365 (M+K), 349 (M+Na), 327 (M+H), 295 (M-OMe)

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 7.56 (s, 1H, Ar), 7.33-7.26 (m, 5H, Ar), 6.91 (s, 1H, Ar), 4.24 (s, 3H, OMe), 4.13 (s, 2H, CH₂Ar), 3.93 (s, 3H, OMe), 3.91 (s, 3H, OMe).

25

Example 9:

6,7-Dimethoxy-2-(2,2-dimethylpropyl)-benzofuran-4-carboxylic acid methyl ester (180 mg, 94%) was obtained as a glassy solid by reacting 6,7-dimethoxy-2-(2,2-dimethylpropanoyl)-benzofuran-4-carboxylic acid methyl ester (200 mg, 0.62 mmol) with sodium cyanoborohydride (392 mg, 6.24 mmol) and trimethylsilyl chloride (789 µl, 6.24 mmol).

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 7.57 (s, 1H, Ar), 6.92 (s, 1H, Ar), 4.29 (s, 3H, OMe), 3.95 (s, 3H, OMe), 3.94 (s, 3H, OMe), 2.66 (s, 2H, CH₂), 1.03 (s, 9H, tert-Bu)

35

General procedure E: Alternative route to obtain benzofuran derivatives IX

Compound **XI** (1 eq.) was reacted with compound **XII** (1 eq.) (obtained by sodium borohydride reduction of the corresponding ketone) as described previously (Wipf, P.; Weiner, W.S. *J. Org. Chem.* 1999, 64, 5321-5324) to give compound **XIII**.

5

Example 10:

3-(1-Cyclopropyl-trimethylsilylprop-2-ynyloxy)-4,5-dimethoxy-benzoic acid methyl ester was obtained as a colorless oil in 60% yield from 1-cyclopropyl-3-(trimethylsilyl)prop-2yn-1-ol. The same compound can also be obtained by a copper-1
10 (CuCl)-catalyzed process as described by Godfrey et al. (Godfrey Jr., J.D. et al, *Tetrahedron Lett.* 1994, 35, 6405-6408).

MS: M⁺ 362

Ether **XIII** (1 eq.), K₂CO₃ (3 eq.) and KF (3 eq.) were dissolved in methanol. The
15 mixture was stirred and the reaction was monitored by TLC. After 25 minutes the reaction was complete and the mixture was partitioned between of water and a mixture of hexane/ethyl acetate 4:1. The layers were separated, the aqueous layer was washed three times with of hexane/ethyl acetate 4:1. The organic layers were combined and evaporated to give the resulting acetylene which was used directly for
20 the next step.

Example 11:

3-(1-Cyclopropyl-prop-2-ynyloxy)-4,5-dimethoxy-benzoic acid methyl ester (17.5 g, 71 %)
25 was obtained as a white solid by crystallization from hexane/ether by reacting 3-(1-cyclopropyl-trimethylsilylprop-2-ynyloxy)-4,5-dimethoxy-benzoic acid methyl ester (30.87 g, 80 mmol) with K₂CO₃ (30 g) and KF (30 g) 17.5 g, 60.4 mmol, 71 %. mp. 93-94 °C.

The acetylene previously obtained (1 eq.) was dissolved in DMF (tech.) containing 5%
30 N,N-diethylaniline and the reaction mixture was stirred for 2h at 150°C. After work-up and crystallisation from ether the compound **IX** was obtained.

Example 12:

6,7-Dimethoxy-2-cyclopropylmethyl-benzofuran-4-carboxylic acid methyl ester (50%)
35 was obtained as white-red crystals by reacting in an intramolecular fashion 3-(1-

cyclopropyl-prop-2-ynyloxy)-4,5-dimethoxy-benzoic acid methyl ester (88 g, 0.316 mol)

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 7.57 (s, 1H, Ar), 7.00 (m, 1H, Ar), 4.28 (s, 3H, OMe), 3.95 (s, 3H, OMe), 3.94 (s, 3H, OMe), 2.70 (d, 2H, J = 6.6), 1.95-1.75 (m, 1H),
5 0.62-0.59 (m, 2H), 0.29-0.25 (m, 2H).

General procedure F: Reduction of the ester to alcohol

Under argon, ester XIV (1 eq.) was dissolved in THF freshly distilled and treated with
10 LiAlH₄ (3 eq.) The reaction was stirred at 60°C for 2h until disappearance of the starting material. The reaction was quenched with 0.5 N HCl, the white precipitate filtrated and washed with ether. The organic layer was dried with MgSO₄ and the solvents evaporated to give the alcohol XV.

15 Example 13:

(6,7-Dimethoxy-2-phenylmethyl-benzofuran-4-yl)-methanol (82 mg, 90%) was obtained by reacting 6,7-dimethoxy-2-phenylmethyl-benzofuran-4-carboxylic acid methyl ester (100 mg, 0.30 mmol) with LiAlH₄ (33 mg, 0.91 mmol).

MS ESI: 321 (M+Na)

20

General procedure G: Oxidation of the alcohol to aldehyde

Under argon, alcohol XV (1 eq.) was dissolved in dichloromethane and MnO₂ (10 eq.) was added. The reaction mixture was stirred 3h at r.t. and filtrated over celite. The celite was washed with excess of CH₂Cl₂, the solvent evaporated to give the aldehyde

25 XVI.

Example 14:

6,7-Dimethoxy-2-phenylmethyl-benzofuran-4-carbaldehyde (119 mg, 62%) was obtained as an oil by reacting (6,7-dimethoxy-2-phenylmethyl-benzofuran-4-yl)-
30 methanol (192 mg, 1.29 mmol) with MnO₂ (1.121 g, 12.90 mmol)

General procedure H: Reduction of the ester to the aldehyde

Under argon at 0°C, to a solution of sodium dihydro-bis(2-methoxyethoxy)-aluminate
35 (Red Al, ca. 3.5 M in toluene) (3.1 eq.) in toluene (over 4Å Mol. sieve) was added dropwise, within 30 min, morpholine (3.4 eq.) in toluene. The resulting solution was

stirred for 30 min, and it was slowly added at -30°C to a solution of the ester **XIV** (1 eq.) in toluene. The reaction mixture was then stirred for 4h at -15°C and quenched by addition of 3N NaOH and stirred until it reached r.t. The reaction mixture was then diluted with ice water and extracted with toluene. The organic layer was washed with
5 water, dried over MgSO_4 and the solvent evaporated to give compound **XVI**.

Example 15:

6,7-Dimethoxy-2-cyclopropylmethyl-benzofuran-4-carbaldehyde (1.16 g, 66%) was obtained as light yellow crystals after crystallisation in 5:1 ethyl acetate/hexane by
10 reacting sodium dihydro-bis(2-methoxyethoxy)-aluminate (Red Al, ca. 3.5M in toluene) (6.5 ml, 22.6 mmol) with morpholine (2.1 ml, 24.0 mmol) and 6,7-dimethoxy-2-cyclopropylmethyl-benzofuran-4-carboxylic acid methyl ester (2.05 g, 7.0 mmol)
NMR CDCl_3 300 MHz δ in ppm J in Hz: 10.06 (s, 1H, COH), 7.32 (s, 1H, Ar) 7.51 (s, 1H, Ar), 4.35 (s, 3H, OMe), 3.97 (s, 3H, OMe), 2.72 (d, 2H, J = 7.1), 1.19-1.17 (m,
15 1H), 0.64-0.60 (m, 2H), 0.29-0.27 (m, 2H).
MP $44-46^{\circ}\text{C}$

Example 16:

6,7-Dimethoxy-2-(2,2dimethylpropyl)-benzofuran-4-carbaldehyde (115 mg, 80%) was
20 obtained as an oil from the reaction of sodium dihydro-bis(2-methoxyethoxy)-aluminate (Red Al, ca. 3.5M in toluene) (500 μl , 1.62 mmol) with morpholine (159 μl , 1.83 mmol) and 6,7-dimethoxy-2-(2,2dimethylpropyl)-benzofuran-4-carboxylic acid methyl ester (160 mg, 0.522 mmol).

25

General procedure I: Coupling with anilinopropionitrile

Under argon at 10°C , the aldehyde **XVI** (1 eq.) was dissolved in DMSO (25 ml) and freshly crystallized 3-anilinopropionitrile (1.1 eq.) was added. Potassium tert-butoxide (1.15 eq.) was added in portions to the reaction mixture. The solution was stirred at
30 10°C for 1h and at r.t. for 3h. The reaction mixture was poured into ice-water and extracted 3 times with ethyl acetate. The organic layer was washed with water and dried over MgSO_4 . The solvents were evaporated to give compound **XVII**.

Example 17:

35 3-Anilino-2-(6,7-dimethoxy-2-cyclopropylmethyl-benzofuran-4-ylmethyl)-acrylonitrile (1.87 g, 49%) was obtained after purification by flash chromatography 2:3 ethyl

acetate/hexane to as a yellow oil from the reaction of 6,7-dimethoxy-2-cyclopropylmethyl-benzofuran-4-carbaldehyde (2.6 g, 9.9 mmol) with 3-anilinopropionitrile (1.44 g, 10.9 mmol) and potassium tert-butoxide (1.28 g, 11.5 mmol)

5

Example 18:

3-Anilino-2-(6,7-dimethoxy-2-phenylmethyl-benzofuran-4-ylmethyl)-acrylonitrile (88 mg, 53%) was obtained after purification by flash chromatography (2:3 ethyl acetate/hexane) as a yellow oil by reacting of 6,7-dimethoxy-2-phenylmethyl-benzofuran-4-carbaldehyde (119 mg, 0.40 mmol) with 3-anilinopropionitrile (58 mg, 0.44 mmol) and potassium tert-butoxide (52 mg, 0.46 mmol)
MS ESI: 423 (M-H)

10

Example 19:

3-Anilino-2-(6,7-dimethoxy-2-(2,2-dimethylpropyl)-benzofuran-4-ylmethyl)-acrylonitrile (91 mg, 54%) was obtained after purification by flash chromatography (2:3 ethyl acetate/hexane) as a yellow oil by reacting 6,7-dimethoxy-2-(2,2dimethylpropyl)-benzofuran-4-carbaldehyde (115 mg, 0.42 mmol) with 3-anilinopropionitrile (60 mg, 0.46 mmol) and potassium tert-butoxide (54 mg, 0.48 mmol)
MS ESI: 427 (M+Na)

15

20

General procedure J: Cyclisation with guanidine

Under argon, to a solution of guanidine hydrochloride (3 eq.) in ethanol potassium tert-butoxide (3 eq.) was added and the mixture was stirred for 15 min. The fine precipitate was filtered through celite under argon and the filtrate was added to a solution of compound XVII (1 eq.) in ethanol. The reaction mixture was stirred under reflux conditions for 8h. After cooling the reaction to r.t., then to -20°C compound I precipitated.

25

Example 20:

5-(2-Cyclopropylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (1.16 g, 96%) was obtained as a white-yellow precipitate by reacting guanidine hydrochloride (1.37 g, 14.44 mmol), potassium tert-butoxide (1.62 g, 14.44 mmol) and 3-anilino-2-(6,7-dimethoxy-2-cyclopropylmethyl-benzofuran-4-ylmethyl)acrylonitrile (1.87 g, 4.81 mmol)

30

35

NMR CD₃OD 500 MHz δ in ppm J in Hz: 6.95-7.35 (m, 1H, Ar), 6.78 (s, 1H, Ar) 6.44 (s, 1H, Ar), 4.03 (s, 3H, OMe), 3.85 (s, 3H, OMe), 3.81 (s, 2H, CH₂), 2.66 (d, 2H, J = 6.8), 1.14-1.04 (m, 1H), 0.59-0.54 (m, 2H), 0.28-0.25 (m, 2H).

MS ISP: 397 (M+Na, 25%), 355 (M+H, 100%).

5 HPLC purity RP C₁₈ Discovery: 98%

Example 21:

5-(2-Benzyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (20 mg, 36%) was obtained as a white-yellow precipitate by reacting guanidine hydrochloride (40 mg, 0.42 mmol), potassium tert-butoxide (47 mg, 0.42 mmol) and 3-anilino-2-(6,7-dimethoxy-2-phenylmethyl-benzofuran-4-ylmethyl)-acrylonitrile (59 mg, 0.14 mmol)

MS ESI: 391 (M+H)

HPLC purity RP C₁₈ Discovery: 97%

15 **Example 22:**

5-(6,7-Dimethoxy-2-(2,2-dimethylpropyl)-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (28 mg, 33%) was obtained as a white-yellow precipitate by reacting guanidine hydrochloride (64 mg, 0.67 mmol), potassium tert-butoxide (75 mg, 0.67 mmol) and 3-anilino-2-(6,7-dimethoxy-2-(2,2-dimethylpropyl)-benzofuran-4-ylmethyl)-acrylonitrile (91 mg, 0.22 mmol)

MS ESI: 371 (M+H)

HPLC purity RP C₁₈ Discovery: 99%

Example 23:

25 Following general procedure C, 2-(biphenyl-4-carboxyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.6 g, 95%) was obtained as a yellow solid by reacting methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (1.0 g, 4.16 mmol) with K₂CO₃ powder (1.44 mg, 10.4 mmol) and 4-phenyl-phenacylbromide (2.29 g, 8.32 mmol)

30 **Example 24:**

Following general procedure C, 6,7-dimethoxy-2-(4-methoxy-benzoyl)-benzofuran-4-carboxylic acid methyl ester (0.331 g, 43%) was obtained as a yellow solid by reacting methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (0.5 g, 2.08 mmol) with K₂CO₃ powder (0.69 mg, 4.9 mmol) and 2-bromo-4-methoxyacetophenone (0.98 g, 4.16 mmol)

NMR CDCl₃ 300 MHz δ in ppm J in Hz: 8.01 (d, 2H, Ar), 7.92 (s, 1H, Ar), 7.64 (s, 1H, Ar), 6.94 (d, 2H, Ar), 4.27 (s, 3H, OMe), 3.93 (s, 3H, OMe), 3.90 (s, 3H, OMe), 3.83 (s, 3H, OMe).

5 **Example 25:**

Following general procedure C, 2-(4-fluoro-benzoyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (2.37, 82%) was obtained as a yellow solid by reacting methyl 2-formyl-3-hydroxy-4,5-dimethoxy-benzoate (2.0 g, 8.32 mmol) with K₂CO₃ powder (2.30 mg, 16.62 mmol) and 2-bromo-4-fluoracetopenon (2.71 g, 12.49 mmol)

10

Example 26:

Following general procedure D, 2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (474 mg, 98%) was obtained as yellow oil by reacting trimethylsilyl chloride (1.52 ml, 12.0 mmol) and sodium cyanoborohydride (754 mg, 12.0 mmol).

15

Example 27:

Following general procedure D, 6,7-dimethoxy-2-(4-methoxy-benzyl)-benzofuran-4-carboxylic acid methyl ester (131 mg, 93%) was obtained as yellow oil by reacting trimethylsilyl chloride (0.341 ml, 2.7 mmol) and sodium cyanoborohydride (169 mg, 2.7 mmol).

20

MS ESI 1: 379 (M+Na)

Example 28:

Following general procedure D, 2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (868 mg, 90%) was obtained as yellow oil by reacting trimethylsilyl chloride (3.64 ml, 28.7mmol) and sodium cyanoborohydride (1.8 g, 28.7 mmol).

25

MS ESI + 329 (M-Me)

30

Example 29: Cleavage of the tBu ester

Under argon at 0°C 6,7-Dimethoxy-benzofuran-2,4-dicarboxylic acid 2-tert-butyl ester 4-methyl ester (10g, 29.7 mmol) was dissolved in TFA (23 ml, 297.3 mmol) and the reaction mixture was stirred for 3h allowing the temperature to reach r.t. The excess of TFA was evaporated and the resulting pink solid was triturated in ether. The resulting solid was dissolved in NaHCO₃ saturated solution, the aqueous layer was

35

extracted with ether then the basic aqueous solution was acidified to pH=1 with 6N HCl and the product was extracted with ethyl acetate. The ethyl acetate organic layer was washed with brine, dried over MgSO₄ and the solvent evaporated to give 6,7-dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (6.43, 77%) as a white solid.

MS ESI - :279 (M-H)

General procedure K: coupling with amines

Under argon at 0°C, to a solution of the acid 6,7-Dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (1 eq.) in CH₂Cl₂/DMF (20:1) were added HOBt (1.2 eq.), EDC(1.2 eq.), the secondary or primary amine (1.1 eq.) and finally triethyl amine (3eq.). The reaction mixture was stirred overnight. Some additional CH₂Cl₂ was added and the organic layer was washed with HCl 1N, H₂O, NaHCO₃ sat and brine. After drying the organic layer over MgSO₄ the solvent was evaporated to give compound IX.

Example 30:

2-Dimethylcarbamoyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.64 g, 97%) was obtained by reacting give 6,7-dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (1.5g, 5.35 mmol) with HOBt (868 mg, 6.42 mmol) EDC (1.23 g, 6.42 mmol), dimethylamine (480 mg, 5.88mmol), and Et₃N (2.23 ml, 16.05 mmol). MS ESI+: 308 (M+H)

Example 31:

2-(3,4-Dihydro-1H-isoquinoline-2-carbonyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (2.09 g, 98%) was obtained by reacting give 6,7-dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (1.5g, 5.35 mmol) with HOBt (868 mg, 6.42 mmol) EDC (1.23 g, 6.42 mmol), tetrahydrosisoquinoline (0.747 ml, 5.88mmol), and Et₃N (2.23 ml, 16.05 mmol). MS ESI+: 396 (M+H)

Example 32:

6,7-Dimethoxy-2-(morpholine-4-carbonyl)-benzofuran-4-carboxylic acid methyl ester (1.85 g, 98%) was obtained by reacting give 6,7-dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (1.5g, 5.35 mmol) with HOBt (868 mg, 6.42 mmol)

EDC (1.23 g, 6.42 mmol), morpholine (0.513 ml, 5.88mmol), and Et₃N (2.23 ml, 16.05 mmol).

MS ESI+: 350 (M+H)

5 **Example 33:**

2-(4-Fluoro-phenylcarbamoyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (978 mg, 52%) was obtained by reacting give 6,7-dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (1.4g, 4.99 mmol) with HOBt (810 mg, 5.99 mmol) EDC (1.15 g, 5.99 mmol), 4-fluoroaniline (0.527 ml, 5.49 mmol), and Et₃N (2.1 ml, 14.98 mmol).

MS ESI-:372 (M+H)

Example 34: Reduction of the acid

Under argon, 6,7-dimethoxy-benzofuran-2,4-dicarboxylic acid 4-methyl ester (66.46 g, 237 mmol) was dissolved in THF dist. and a 2M solution of BH₃.DMS (116 ml, 0.260 mmol) was slowly added. The reaction mixture was stirred at reflux overnight. After cooling to r.t the reaction was quenched with MeOH. The solvents were then evaporated and the resulting oil was dissolved in ethyl acetate. The organic layer was washed with NaHCO₃ sat., H₂O and brine, dried over MgSO₄. 2-Hydroxymethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (34 g, 54%) was obtained as yellow-white solid after purification by FC Hexane/ethyl acetate (3:2).

MS ESI+:267 (M+H)

25 **General Procedure L: Coupling with halogen derivatives**

Under argon, to a solution of 2-hydroxymethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid (1 eq.) in DMF dried over molecular sieve was added portionwise NaH 60% (1.1 eq.). Halogen derivatives XR⁹ (1.05 eq.) were slowly added to the solution. The reaction was stirred 1h at r.t and was quenched with some H₂O. After evaporation of the DMF the resulting oil was partitioned between ethyl acetate and NaHCO₃. After extraction of the water layer with ethyl acetate the organic layer was washed with H₂O, KHSO₄ 1N and brine, dried over MgSO₄ and the solvents evaporated. The pure compound IX was obtained after purification by FC hexane/ethyl acetate (4:1).

Example 35:

2-Allyloxymethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (10.43 g, 60%) was obtained as a yellow oil by reacting 2-hydroxymethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid (15g, 56.39 mmol) with NaH 60% (2.48 g, 62.03 mmol) and allyl bromide (5.01 ml, 59.21 mmol).
LC MS clean but no ionisation

Example 36:

6,7-Dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-carboxylic acid methyl ester (480 mg, 40%) was obtained as a crystalline solid by reacting 2-hydroxymethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid (840 mg, 3.15 mmol) with NaH 60% (139 mg, 3.47 mmol), K₂CO₃ (436 mg, 3.15 mmol) and methoxybenzyl chloride (0.449 ml, 3.31 mmol).
NMR CDCl₃ 300 MHz δ in ppm J in Hz: 7.61 (s, 1H, Ar), 7.29 (d, 2H, Ar), 7.21 (s, 1H, Ar), 6.28 (d, 2H, Ar), 4.61 (s, 2H, CH₂), 4.56 (s, 2H, CH₂), 4.28 (s, 3H, OMe), 3.93 (s, 6H, 2xOMe), 3.80 (s, 3H, OMe).

Example 37:

Following general procedure F, (2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-yl)-methanol (395 mg, 90%) was obtained as a transparent oil by reacting 2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (474 mg, 1.18 mmol) with LiAlH₄ (135 mg, 3.53 mmol).
MS ESI +: 357 (M-OH)

Example 38:

Following general procedure F, [6,7-dimethoxy-2-(4-methoxy-benzyl)-benzofuran-4-yl]-methanol (79 mg, 66%) was obtained as an oil by reacting 6,7-dimethoxy-2-(4-methoxy-benzyl)-benzofuran-4-carboxylic acid methyl ester (131 mg, 0.36 mmol) with LiAlH₄ (28 mg, 0.73 mmol).
MS ESI +: 311 (M-OH)

Example 39:

Following general procedure F, [2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-yl]-methanol (709 mg, 86%) was obtained as an oil by reacting 2-(4-fluoro-benzyl)-6,7-

dimethoxy-benzofuran-4-carboxylic acid methyl ester (868 mg, 2.60 mmol) with LiAlH_4 (197 mg, 5.21 mmol).

MS ESI +: 299 (M-OH)

5 **Example 40:**

Following general procedure F, (2-dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-yl)-methanol (507 g, 60%) was obtained as an oil by reacting 2-dimethylcarbamoyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.0 g, 2.60 mmol) with LiAlH_4 (494 mg, 13.01 mmol).

10 MS ESI +: 266 (M+H)

Example 41:

Following general procedure F, [2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-yl]-methanol (970 mg, 77%) was obtained as an oil by reacting

15 2-(3,4-dihydro-1H-isoquinoline-2-carbonyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.4 g, 3.54 mmol) with LiAlH_4 (537 mg, 14.16 mmol).

MS ESI +: 354 (M+H)

Example 42:

20 Following general procedure F, (6,7-dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-yl)-methanol (800 mg, 60%) was obtained as an oil by reacting 6,7-dimethoxy-2-(morpholine-4-carbonyl)-benzofuran-4-carboxylic acid methyl ester (1.52 g, 4.35 mmol) with LiAlH_4 (660 mg, 17.40 mmol).

MS ESI +: 308 (M+H)

25

Example 43:

Following general procedure F, {2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-yl}-methanol (724 mg, 84%) was obtained as an oil by reacting 2-(4-fluoro-phenylcarbamoyl)-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester

30 (974 mg, 2.6 mmol) with LiAlH_4 (495 mg, 13.0 mmol).

MS ESI +: 330 (M+H) very weak signal

Example 44:

Following general procedure F, (2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-yl)-methanol (1.76 g, 98%) was obtained as an oil by reacting 2-allyloxymethyl-6,7-

35

dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.95 g, 6.36 mmol) with LiAlH_4 (483 mg, 12.73 mmol).

MS ESI+: 261 (M-OH)

5 **Example 45:**

Following general procedure F, [6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-yl]-methanol (390 mg, 95%) was obtained as an oil by reacting 6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-carboxylic acid methyl ester (445 mg, 1.15 mmol) with LiAlH_4 (87 mg, 2.30 mmol).

10 MS ESI+: 341 (M-OH)

Example 46:

Following general procedure F, (6,7-dimethoxy-2-phenyl-benzofuran-4-yl)-methanol (150 mg, 74%) was obtained as an oil by reacting 6,7-dimethoxy-2-phenylbenzofuran-4-carboxylic acid methyl ester (235 mg, 0.735 mmol) with LiAlH_4 (14 mg, 0.377 mmol). 6,7-dimethoxy-2-phenylbenzofuran-4-carboxylic acid methyl ester (240 mg, 22%) was obtained following the palladium coupling procedure [J. Org. Chem. 1996, 61, 9280-9288; Bioorg. Med. Chem., 1999, 7, 6, 1131-1140] by reacting 2-bromo-3-hydroxy-4,5-dimethoxy-benzoic acid methyl ester (1 g, 3.44 mmol) (P. Wipf and W. S. Weiner, J. Org. Chem. 1999, 64, 5321-5324) with phenylacetylen (525 mg, 5.15 mmol), CuI (13 mg, 0.07 mmol) and $\text{Pd(PPh}_3\text{)Cl}_2$ (24 mg, 0.04 mmol) in $\text{Et}_3\text{N/DMF}$ (3:1).

MS ESI+: 267 (M-OH)

25 **Example 47:**

Following general procedure F, [6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-yl]-methanol (3.94 g, 98%) was obtained as an oil by 6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-carboxylic acid methyl ester (4.29 g, 11.16 mmol) with LiAlH_4 (980 mg, 24.5 mmol).

30 MS ESI+: 357 (M+H)

Example 48:

Following general procedure G, 2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-carbaldehyde (105 mg, 27%) was obtained as a solid by reacting (2-biphenyl-4-

ylmethyl-6,7-dimethoxy-benzofuran-4-yl)-methanol (395 mg, 1.06 mmol) with MnO₂ (1.73 g, 19.9 mmol)

MS ESI +: 373 (M+H)

5 Example 49:

Following general procedure G, 6,7-dimethoxy-2-(4-methoxy-benzyl)-benzofuran-4-carbaldehyde (20 mg, 25%) was obtained as an oil by reacting [6,7-dimethoxy-2-(4-methoxy-benzyl)-benzofuran-4-yl]-methanol (79 mg, 0.24 mmol) with MnO₂ (210 mg, 2.42 mmol)

10 MS ESI +: 329 (M+H)

Example 50:

Following general procedure G, 2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-carbaldehyde (396 mg, 56%) was obtained as an oil by reacting [2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-yl]-methanol (709 mg, 2.24 mmol) with MnO₂ (1.948 g, 22.41 mmol)

15 MS ESI +: 315 (M+H)

Example 51:

20 Following general procedure G, 2-dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-carbaldehyde (496 mg, 77%) was obtained as a brown oil by reacting [2-dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-yl]-methanol (500 mg, 1.88 mmol) with MnO₂ (1.64 g, 18.84 mmol)

MS ESI +: 264 (M+H)

25

Example 52:

Following general procedure G, 2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-carbaldehyde (638 mg, 71%) was obtained as a brown oil by reacting [2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-yl]-

30 methanol (907 mg, 2.56 mmol) with MnO₂ (2.23 g, 25.66 mmol)

MS ESI +: 352 (M+H)

Example 53:

Following general procedure G, 6,7-Dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-carbaldehyde (489 mg, 64%) was obtained as an orange oil by reacting (6,7-

35

dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-yl)-methanol (765 mg, 2.49 mmol)
with MnO₂ (2.16 g, 24.89 mmol)
MS ESI +: 306 (M+H)

5 **Example 54:**

Following general procedure G, 2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-carbaldehyde (197 mg, 28%) was obtained as an oil by reacting {2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-yl}-methanol (718 mg, 2.16 mmol) with MnO₂ (1.88 g, 21.6 mmol)

10 MS ESI +: 330 (M+H)

Example 55:

Following general procedure G, 2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-carbaldehyde (2.01 g, 72%) was obtained as an oil by reacting (2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-yl)-methanol (2.82, 10.17 mmol) with MnO₂ (8.80 g, 101.69 mmol)

15 MS ESI +: 277 (M+H)

Example 56:

20 Following general procedure G, 6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-carbaldehyde (244 mg, 72%) was obtained as an oil solid by reacting [6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-yl]-methanol (341 mg, 0.95 mmol) with MnO₂ (0.827 g, 9.51 mmol)

MS ESI +: 357 (M+H)

25

Example 57:

Following general procedure G, 6,7-dimethoxy-2-phenyl-benzofuran-4-carbaldehyde (121 mg, 81%) was obtained as an oil by reacting (6,7-dimethoxy-2-phenyl-benzofuran-4-yl)-methanol (150 mg, 0.53 mmol) with MnO₂ (467 mg, 5.33 mmol)

30 MS ESI +: 283 (M+H)

Example 58:

Following general procedure G, 6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-carbaldehyde (3.50 g, 97%) was obtained as a oil by reacting ([6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-yl]-methanol (3.65 mg, 10.24
5 mmol) with MnO₂ (9.8 g, 102 mmol)
MS ESI +: 355 (M+H)

Example 59:

Following general procedure I, 2-(2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl-3-phenylamino-acrylonitrile (58 mg, 41%) was obtained after purification by
10 flash chromatography 2:3 ethyl acetate/hexane as a yellow oil from the reaction of 2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-carbaldehyde (105 mg, 0.28 mmol) with 3-anilinopropionitrile (45 mg, 0.30 mmol) and potassium tert-butoxide (36.4 mg, 0.32 mmol)

15

Example 60:

Following general procedure I, 2-[6,7-dimethoxy-2-(4-methoxy-benzoyl)-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile which was used for the next step without further purification, was obtained from the reaction 6,7-dimethoxy-2-(4-methoxy-benzyl)-benzofuran-4-carbaldehyde (20 mg, 0.06 mmol) with 3-anilinopropionitrile (9
20 mg, 0.06 mmol) and potassium tert-butoxide (8 mg, 0.07 mmol)

Example 61:

Following general procedure I, 2-[2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile which was used for the next step without further
25 purification, was obtained from the reaction 2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-carbaldehyde (395 mg, 1.25 mmol) with 3-anilinopropionitrile (200 mg, 1.38 mmol) and potassium tert-butoxide (170 mg, 1.50 mmol)
MS ESI -: 441 (M-H)

30

Example 62:

Following general procedure I, 2-(2-dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile which was used for the next step without further purification, was obtained from the 2-dimethylaminomethyl-6,7-dimethoxy-
5 benzofuran-4-carbaldehyde (350 mg, 1.33 mmol) with 3-anilinopropionitrile (223 mg, 1.53 mmol) and potassium tert-butoxide (179 mg, 1.60 mmol)
MS ESI -: 390 (M-H)

Example 63:

10 Following general procedure I, 2-[2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile (414 mg, 52%) was obtained after purification by flash chromatography 2:3 ethyl acetate/hexane as a yellow oil from the reaction of 2-2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-carbaldehyde (584 mg, 1.66 mmol) with 3-anilinopropionitrile
15 (279 mg, 1.91 mmol) and potassium tert-butoxide (224 mg, 1.99 mmol)
MS ESI +: 480 (M+H)

Example 64:

Following general procedure I, 2-(6,7-dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile (348 mg, 50%) was obtained after purification
20 by flash chromatography 2:3 ethyl acetate/hexane as a yellow oil from the reaction of 6,7-dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-carbaldehyde (485 mg, 1.59 mmol) with 3-anilinopropionitrile (267 mg, 1.82 mmol) and potassium tert-butoxide (214 mg, 1.90 mmol)
25 MS ESI +: 434 (M+H)

Example 65:

Following general procedure I, 2-{2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-ylmethyl}-3-phenylamino-acrylonitrile which was used for the next step
30 without further purification, was obtained from the 2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-carbaldehyde (197 mg, 0.60 mmol) with 3-anilinopropionitrile (100 mg, 0.69 mmol) and potassium tert-butoxide (80 mg, 0.72 mmol)

Example 66:

Following general procedure I, 2-[6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile (123 mg, 38%) was obtained after purification by flash chromatography 2:3 ethyl acetate/hexane as a yellow oil from the reaction of 6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-carbaldehyde (236 mg, 0.66 mmol) with 3-anilinopropionitrile (106 mg, 0.78 mmol) and potassium tert-butoxide (85 mg, 0.761 mmol)

MS ESI -: 472 (M-H)

Example 67:

Following general procedure I, 2-(2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile (501 mg, 54%) was obtained after purification by flash chromatography 2:3 ethyl acetate/hexane as a yellow oil from the reaction of 2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-carbaldehyde (626 mg, 2.26 mmol) with 3-anilinopropionitrile (364 mg, 2.49 mmol) and potassium tert-butoxide (292 mg, 2.60 mmol)

MS ESI -: 403 (M-H)

Example 68:

Following general procedure I, 2-(6,7-dimethoxy-2-phenyl-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile which was used for the next step without further purification, was obtained from the 6,7-dimethoxy-2-phenyl-benzofuran-4-carbaldehyde (120 mg, 0.426 mmol) with 3-anilinopropionitrile (62 mg, 0.426 mmol) and potassium tert-butoxide (48 mg, 0.426 mmol)

25

Example 69:

Following general procedure I, 3-[6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-yl]-2-phenylaminomethyl-acrylonitrile which was used for the next step without further purification, was obtained from the 6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-carbaldehyde (2.35 g, 6.63 mmol) with 3-anilinopropionitrile (1.02 g, 6.97 mmol) and potassium tert-butoxide (0.86 g, 7.62 mmol)

MS ESI +: 483 (M+H)

30

Example 70:

- 5-(2-Biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine formic acid salt (1 mg) was obtained as a white-yellow solid after purification by prep HPLC (RP C18, 10 mM Formic acid/ACN) by reacting guanidine hydrochloride (33 mg, 0.344 mmol), potassium tert-butoxide (41 mg, 0.367 mmol) and 2-(2-biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl-3-phenylamino-acrylonitrile (57 mg, 0.115 mmol).
MS ESI +: 467 (M+H)

Example 71:

- Following General Procedure J, 5-[6,7-dimethoxy-2-(4-methoxy-benzyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine (20 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (17 mg, 0.18 mmol), potassium tert-butoxide (20 mg, 0.18 mmol) and 2-[6,7-dimethoxy-2-(4-methoxy-benzoyl)-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile.
MS ESI +: 421 (M+H)

Example 72:

- Following General Procedure J, 5-[2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine hydrochloride salt (60 mg) was obtained as a dark yellow solid after crystallisation of the salt in ether/HCl by reacting guanidine hydrochloride (344 mg, 3.60 mmol), potassium tert-butoxide (404 mg, 3.60 mmol) and 2-[2-(4-fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile.
MS ESI +: 409 (M+H)

Example 73:

- Following General Procedure J, 5-(2-dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (310 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (366 mg, 3.83 mmol), potassium tert-butoxide (430 mg, 3.83 mmol) and 2-(2-dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile.
MS ESI +: 358 (M+H)

Example 74:

Following General Procedure J, 5-[2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (234 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (245 mg, 2.56 mmol), potassium tert-butoxide (288 mg, 2.56 mmol) and 2-[2-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile (410 mg, 0.85 mmol)
MS ESI +: 446 (M+H)

Example 75:

Following General Procedure J, 5-(6,7-dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (177 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (228 mg, 2.39 mmol), potassium tert-butoxide (268 mg, 2.39 mmol) and 2-(6,7-dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile (345 mg, 0.79 mmol).
MS ESI +: 400 (M+H)

Example 76:

Following General Procedure J, 5-{2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-ylmethyl}-pyrimidine-2,4-diamine (4 mg) was obtained as a yellow solid after purification by TLC preparative (SiO₂, CH₂Cl₂/MeOH, 9:1) by reacting guanidine hydrochloride (163 mg, 1.05 mmol), potassium tert-butoxide (118 mg, 1.05 mmol) and 2-{2-[(4-fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-ylmethyl}-3-phenylamino-acrylonitrile
MS ESI +: 424 (M+H)

Example 77:

Following General Procedure J, 5-[6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (32 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (99 mg, 0.68 mmol), potassium tert-butoxide (76 mg, 0.68 mmol) and 2-[6,7-dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-ylmethyl]-3-phenylamino-acrylonitrile (110 mg, 0.23 mmol)
MS ESI +: 451 (M+H)

Example 78:

Following General Procedure J, 5-(2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (299 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (531 mg, 3.63 mmol),
5 potassium tert-butoxide (408 mg, 0.68 mmol) and 2-(2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile (490 mg, 1.21 mmol)
MS ESI +: 371 (M+H)

Example 79:

10 Following General Procedure J, 5-(6,7-dimethoxy-2-phenyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (4 mg) was obtained as a yellow solid after crystallisation from ethanol by reacting guanidine hydrochloride (122 mg, 1.28 mmol), potassium tert-butoxide (143 mg, 1.28 mmol) and 2-(6,7-dimethoxy-2-phenyl-benzofuran-4-ylmethyl)-3-phenylamino-acrylonitrile.
15 MS ESI +: 377 (M+H)

Example 80:

Following General Procedure J, 5-[6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (440 mg) was obtained as a yellow
20 solid after crystallisation from ethanol by reacting guanidine hydrochloride (890 mg, 9.3 mmol), potassium tert-butoxide (1.04 g, 9.3 mmol) and 3-[6,7-dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-yl]-2-phenylaminomethyl-acrylonitrile (1.5 g crude).
MS ESI +: 449 (M+H)

25

Example 81:

[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-phenyl-methanone; hydrochloride (107 mg, 25%) was obtained as a yellow solid after crystallisation from ethanol by reacting a solution of 5-[6,7-dimethoxy-2-(2-phenyl-
30 [1,3]dioxolan-2-yl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (440 mg, 0.981 mmol) in ethanol with 6 N HCl for 16h at r.t.
MS ESI +: 405 (M+H)

Example 82:

35 [4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-phenyl-methanol (10 mg) was obtained by reacting [4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-

dimethoxy-benzofuran-2-yl]-phenyl-methanone (20 mg) with NaBH₄. Then the reaction was quenched with 1N HCl, the impurities were extracted with ethyl acetate and the solution neutralised with Na₂CO₃ to pH=7 and saturated with NaOAc. The compound was extracted with ACN dried over MgSO₄ and the solvent evaporated to give the
5 desired compound.
MS ESI +: 407 (M+H)

Example 83: Deprotection of hydroxyl

Under argon, to a solution of 5-(2-allyloxymethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (610 mg, 1.64 mmol) in ACN was added Pd(PPh₃)₄
10 complex (571 mg, 0.424 mmol) and ammonium formate (519 mg, 8.23 mmol). The reaction mixture was stirred at reflux for 2h. Then the reaction was quenched with 1N HCl, the impurities were extracted with ethyl acetate and the solution neutralised with Na₂CO₃ to pH=7 and saturated with NaOAc. The compound was extracted with ACN
15 dried over MgSO₄ and the solvent evaporated to give [4-(2,4-diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-methanol (497 mg, 84%) as a yellow solid.

General Procedure M: Protection of the keto group

20 Under argon, to a solution of compounds VI (1 eq.) in dry toluene was added ethylene glycol (6.5 eq.) and p-toluene sulfonic acid (0.15 eq.). The reaction mixture was stirred at reflux for 16h and the water produce during the reaction was distilled using a Dean-stark trap. The reaction mixture was diluted with ether and the organic layer was washed with H₂O, brine and dried over MgSO₄. Purification by FC hexane/ethyl
25 acetate (19:1) gave the expected compound X.

Example 84:

6,7-Dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-carboxylic acid methyl ester (5.08 g, 62%) was obtained as an oil by reacting 2-benzoyl-6,7-dimethoxy-
30 benzofuran-4-carboxylic acid methyl ester (7.22g, 21.21 mmol) with ethylene glycol (7.71 ml, 138.5 mmol) and pTsOH (609 mg, 3.2 mmol)

General Procedure N: Transformation to chloro derivatives

35 Under argon, to a solution of the alcohol (1eq.) in ether or CHCl₃ was added thionyl chloride (1.5-1.7 eq.) and the reaction mixture was stirred for 1h at reflux. In the case

of ether as solvent the organic layer could be washed with H₂O, dried over MgSO₄ and the solvent evaporated to give the chloride derivative VIII.2. In the case of CHCl₃ as solvent, the solvent and the excess of thionyl chloride were distilled under reduced pressure to give the chloride derivative XIX.

5

Example 85:

2-Chloromethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.04 g, 97%) was obtained as a yellow solid by reacting 2-hydroxymethyl-6,7-dimethoxy-benzofuran-4-carboxylic acid methyl ester (1.0 g, 3.75 mmol) with thionyl chloride
10 (0.460 ml, 6.38 mmol) and pyridine (0.04 ml, cat.).

Example 86:

[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-methanol (200 mg, 0.60 mmol) was reacted with thionyl chloride (0.066 ml, 0.90 mmol) to give after
15 direct evaporation of the solvent 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine as a yellow oily compound which was used in the next step without further purification.

20 General Procedure O: Friedel Craft's alkylation

Under argon at 50°C, to a solution of the aromatic derivatives R⁵ (10 eq.) in CHCl₃ was added ZnCl₂ (1.2-2.4 eq.). A solution of the chloride derivative (1 eq.) in CHCl₃ was then slowly added within 30 min. After stirring at 50°C for 1-2h the reaction was quenched with H₂O and the reaction mixture was partitioned between CH₂Cl₂ and
25 NaHCO₃ sat. The organic layer was separated, dried over MgSO₄ and the solvent evaporated. The pure derivatives IX were obtained after purification by FC hexane/ethyl acetate (3:1). The pure derivatives I were obtained after dissolution the crude mixture in 1N HCl, extraction of the nonpolar impurities with ether, basification with Na₂CO₃ to pH=8 and extraction of the compound with ethyl acetate or CHCl₃.
30 The organic layer was then dried over MgSO₄ and the solvent evaporated to give the pure compound I.

Example 87:

5-[2-(3,4-Dimethoxy-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (45 mg) was obtained as a dark yellow solid by reacting 5-(2-chloromethyl)-
35

6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (155 mg, 0.454 mmol) with veratrol (0.529 ml, 4.45 mmol) and ZnCl_2 (68 mg, 0.50 mmol)

MS ESI +: 451 (M+H)

Example 88:

- 5 4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-phenol (3 mg) was obtained after purification of a part of the reaction mixture by TLC prep (SiO_2) as a white-yellow solid by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (50 mg, 0.143 mmol) with phenol (135 mg, 1.43 mmol) and ZnCl_2 (23 mg, 0.172 mmol).

- 10 MS ESI +: 407 (M+H)

Example 89:

- 5-[6,7-Dimethoxy-2-(1-methyl-1H-pyrrol-2-ylmethyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (3 mg) was as a yellow solid by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (50 mg, 0.143 mmol) with methylpyrrole (0.128 ml, 1.43 mmol) and ZnCl_2 (47 mg, 0.34 mmol).
MS ESI +: 394 (M+H)

20 **General procedure P: Coupling**

- Under argon, to a solution of pyrrol, indol, thiophenol or phenol HXR^{10} (1.5-2 eq.) in DMF was added NaH (1.5-2 eq.) and the mixture was stirred for 1 h. 5-(2-Chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (1 eq.) in DMF was then slowly added. The reaction mixture was stirred for 18h at r.t. After
25 quenching the reaction with H_2O , the mixture was partitioned between NaHCO_3 sat. and CHCl_3 . The aqueous layer was extracted several times with CHCl_3 dried over MgSO_4 and the solvent was evaporated. The residue was dissolved in 1N HCl, extracted with ether, basified with Na_2CO_3 to pH=8 and extracted with CHCl_3 . The CHCl_3 layer was dried over MgSO_4 , the solvent evaporated to give the pure
30 compounds I. The salts were obtained after treating the free base with the corresponding acidic aqueous solution and freeze-drying.

Example 90:

- 5-[2-(4-Fluoro-phenoxy-methyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine; formic acid salt (48 mg, 76%) was obtained by reacting 5-(2-chloromethyl-

6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (52 mg, 0.151 mmol), NaH (9 mg, 0.226 mmol), and 4-fluorophenol (25 mg, 0.226 mmol).

MS ESI +: 471 (M+H)

5 **Example 91:**

5-[6,7-Dimethoxy-2-(4-trifluoromethyl-phenoxy)methyl]-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine; formic acid salt (33 mg, 47%) was obtained by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (52 mg, 0.151 mmol), NaH (9 mg, 0.226 mmol), and 4-trifluorophenol (337 mg, 0.226 mmol).

10 MS ESI +: 521 (M+H)

1 **Example 92:**

5-[2-(4-Fluoro-phenylsulfanylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine; formic acid salt (15 mg) was obtained by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (52 mg, 0.151 mmol), NaH (12 mg, 0.303 mmol), and 4-fluorothiophenol (0.032 ml, 0.303 mmol).

MS ESI+. 487 (M+H)

20 **Example 93:**

N-{4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethylsulfanyl]-phenyl}-acetamide; formic acid salt (20 mg) was obtained by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (52 mg, 0.151 mmol), NaH (12 mg, 0.303 mmol), and 4-acetaminothiophenol (56 mg, 0.303 mmol).

25 MS ESI+ : 526 (M+H)

Example 94:

4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethoxy]-phenol (3 mg) was obtained after purification of a part of the reaction mixture by TLC preparative (SiO₂) by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (75 mg, 0.215 mmol), NaH (13 mg, 0.322 mmol), and 4-hydroxyphenol (38 mg, 0.344 mmol).

MS ESI+ : 423 (M+H)

Example 95:

5-[2-(4-Amino-phenoxy-methyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine (43 mg) was obtained by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (75 mg, 0.215 mmol), NaH (13 mg, 0.322 mmol), and 4-aminophenol (50 mg, 0.344 mmol).

MS ESI+ : 422 (M+H)

Example 96:

5-(6,7-Dimethoxy-2-pyrrol-1-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (62 mg, 74%) was obtained by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (77 mg, 0.220 mmol), NaH (13 mg, 0.322 mmol), and pyrrole (0.022 ml, 0.322 mmol).

MS ESI+ : 380 (M+H)

Example 97:

5-(2-Imidazol-1-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (24 mg, 28%) was obtained by reacting 5-(2-chloromethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine (77 mg, 0.220 mmol), NaH (13 mg, 0.322 mmol), and imidazol (22 mg, 0.322 mmol).

MS ESI+ : 381 (M+H)

Example 98:

Following procedure C, 5-(2,4-diamino-pyrimidin-5-ylmethyl)-2,3-dimethoxy-phenol (Helv. Chim. Acta 1970, 53, 945) (200 mg, 0.687 mmol) was reacted with K₂CO₃ (71 mg, 0.515 mmol) and ethyl-2-chloroacetoacetate (0.096 ml, 0.687 mmol) to give after work-up using CHCl₃ and NaHCO₃ sat solution and crystallisation in ethanol, 4-(2,4-diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid ethyl ester as a yellow solid (150 mg, 57%)

MS ESI +: 387 (M+H)

Example 99:

4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid was (10mg) obtained as a yellow solid after treating 4-(2,4-diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid ethyl ester with a 2N NaOH solution neutralisation of the solution with HCl and freeze-drying.

MS ESI +: 359 (M+H)

Example 100:

Following General Procedure G, 4-(2,4-diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-carbaldehyde (10 mg) was obtained by reacting [4-(2,4-diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-methanol (15 mg, 0.04 mmol) and MnO₂ (4 mg, 0.4 mmol)

MS ESI+:329 (M+H)

Example 101: Biological results

Antimicrobial susceptibility testing was performed in accordance with the National Committee for Clinical Laboratory Standards (NCCLS) procedure [M7-A5, 2001].

M7-A5 (2001) : Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard —Fifth Edition American National Standard

15

***In vitro* Antibacterial Activity of Compounds**
(Minimum Inhibitory Concentration (MIC) in micrograms/ml)


Microorganism	End Product of Example			Trimethoprim
	20	21	72	
<i>Escherichia coli</i> ATCC 25922	4.0	2.0	2.0	0.25
<i>Staphylococcus aureus</i> ATCC 25923	0.062	0.062	0.25	2
<i>Streptococcus pneumoniae</i> ATCC 49619	0.125	0.50	0.5	2
<i>Staphylococcus epidermidis</i> ATCC 12228	0.062	0.062	0.015	0.062
<i>Enterococcus faecalis</i> ATCC 29212	0.500	0.125	1.0	0.5
<i>Bacillus subtilis</i> ATCC 6051	0.062	0.062	0.25	0.5

Claims

1. Compounds of the general formula I

5

Formula I



10 wherein

R¹ represents straight or branched chain lower alkyl with 2 to 6 carbon atoms; cycloalkylmethyl with 3 to 6 carbon atoms; aryl; arylmethyl or heteroarylmethyl, the aryl and heteroaryl group may be mono-, di- or tri- substituted with halogen, amino, hydroxy, nitro, trifluoromethyl, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; straight or branched chain lower alkylcarbonyl with up to 6 carbon atoms; cycloalkylcarbonyl with 3 to 6 carbon atoms; cycloalkylhydroxymethyl with 3 to 6 carbon atoms; arylcarbonyl, the aryl group may be mono-, di or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; arylhydroxymethyl, the aryl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; straight or branched chain lower alkenyl with 2 to 6 carbon atoms; hydroxy-lower alkyl with 1 to 6 carbon atoms; fluoro-lower alkyl with 1 to 6 carbon atoms; aryloxy-lower alkyl whereby the aryl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; arylthio-lower alkyl whereby

the aryl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; arylamino-lower alkyl whereby the aryl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; lower alkenyloxy lower alkyl whereby the lower alkenyl group may contain 2 to 4 carbon atoms and the lower alkyl group may contain 1 or 2 carbon atoms; benzyloxy lower alkyl whereby the benzyl group may be mono-, di- or tri- substituted with halogen, amino, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different; lower alkylamino lower alkyl whereby the lower alkyl groups may contain 1 to 3 carbon atoms; heterocyclymethyl containing one to three hetero atoms which can be the same or different and which may be substituted with lower alkyl, halogen, amino, lower alkyloxy, hydroxy, lower alkylcarbonylamino, arylcarbonylamino and benzofused derivatives thereof.

R^2 and R^3 independently represent hydrogen; lower alkyl with 1 to 3 carbon atoms; or together a lower alkylene group with 1 to 3 carbon atoms bridging the oxygen atoms and forming a five, six or seven membered ring;

R^4 represents hydrogen; straight or branched chain lower alkyl with 1 to 4 carbon atoms;

and pharmaceutically acceptable salts and N-oxides thereof.

2. Compounds of formula I, wherein R^1 is straight or branched chain lower alkyl with 2 to 6 carbon atoms; cycloalkylmethyl with 3 to 6 carbon atoms; aryl; arylmethyl or heteroarylmethyl, the aryl and heteroaryl group may be mono- or di substituted with halogen, amino, hydroxy, nitro, trifluoromethyl, lower alkyloxy, lower alkylcarbonylamino, arylcarbonylamino, whereby these substituents may be the same or different, and R^2 and R^3 are methyl or together are a methylene group bridging the oxygen atoms to which they are attached and R^4 is hydrogen or methyl.

3. Compounds of formula I, wherein R^1 is 4-methoxy-benzyl, phenyl, benzyl, cyclopropylmethyl, 4-fluoro-benzyl, 3,4-dihydro-1H-isoquinolin-2-ylmethyl, 4-methoxy-benzyloxymethyl, 4-acetylaminophenyl-sulfanyl-methyl, 4-trifluoromethyl-phenoxy-methyl, 4-amino-phenoxy-methyl, allyloxymethyl, phenyl-[1,3]dioxolan-2-yl,

pyrrol-1-ylmethyl, 3,4-dimethoxy-benzyl, 4-hydroxyphenyloxymethyl, phenylcarbonylmethyl, 4-fluoro-phenoxyethyl, 2,2-dimethyl-propyl, 4-fluoro-phenylsulfanylmethyl, hydroxymethyl, formyl, 4-fluoro-phenylamino-methyl, imidazol-1-ylmethyl, dimethylaminomethyl, morpholin-4-ylmethyl, biphenyl-4-ylmethyl, 5 ethoxycarbonyl, carboxy, 4-hydroxybenzyl, furan-2-ylmethyl, thiophen-2-ylmethyl, thiophen-3-ylmethyl, 1-methyl-pyrrol-2-ylmethyl, pyridin-2-ylmethyl, pyridin-3-ylmethyl, pyridin-4-ylmethyl, 4-acetylaminobenzyl, 4-amino-benzyl, 4-nitro-benzyl, 4-acetyl-amino-3-methoxy-benzyl, 4-amino-3-methoxy-benzyl, 3-acetyl-amino-4-methoxy-benzyl, 3-amino-4-methoxy-benzyl, allyl, isopropenyl or halogenmethyl, and R² and R³ 10 are methyl or together are a methylene group bridging the oxygen atoms to which they are attached and R⁴ is hydrogen or methyl.

4. Compounds of formula I, wherein R¹ is 4-methoxy-benzyl, phenyl, benzyl, cyclopropylmethyl, 4-fluoro-benzyl, 3,4-dihydro-1H-isoquinolin-2-ylmethyl, 4-methoxy- 15 benzyloxymethyl, 4-acetylaminophenyl-sulfanyl-methyl, 4-trifluoromethyl-phenoxyethyl, 4-amino-phenoxyethyl, allyloxymethyl, 3,4-dimethoxy-benzyl, 4-hydroxybenzyl, pyridin-2-ylmethyl, pyridin-3-ylmethyl, pyridin-4-ylmethyl, 4-acetylaminobenzyl, 4-amino-benzyl, 4-acetyl-amino-3-methoxy-benzyl, 4-amino-3-methoxy-benzyl, 3-acetyl-amino-4-methoxy-benzyl, 3-amino-4-methoxy-benzyl or aryl 20 methyl and R² and R³ are both methyl and R⁴ is hydrogen.

5. Compounds of formula I, wherein R¹ is 4-methoxy-benzyl, benzyl, cyclopropylmethyl, 4-fluoro-benzyl, 3,4-dihydro-1H-isoquinolin-2-ylmethyl, 4-amino-phenoxyethyl, 4-hydroxybenzyl, pyridin-2-ylmethyl, pyridin-3-ylmethyl, pyridin-4- 25 ylmethyl, 4-acetylaminobenzyl, 4-amino-benzyl, 4-acetyl-amino-3-methoxy-benzyl, 4-amino-3-methoxy-benzyl, 3-acetyl-amino-4-methoxy-benzyl, 3-amino-4-methoxy-benzyl and R² and R³ are both methyl and R⁴ is hydrogen

6. The compounds according to any one of claims 1 to 5

30

5-[6,7-Dimethoxy-2-(4-methoxy-benzyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,

5-(6,7-Dimethoxy-2-phenyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,

35 5-(2-Benzyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,

- 5-(2-Cyclopropylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-[2-(4-Fluoro-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5
- 5-[2-(3,4-Dihydro-1H-isoquinolin-2-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 10
- 5-[6,7-Dimethoxy-2-(4-methoxy-benzyloxymethyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 15
- 5-[6,7-Dimethoxy-2-(4-trifluoromethyl-phenoxy-methyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 20
- 5-[2-(4-Amino-phenoxy-methyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 25
- 5-(6,7-Dimethoxy-2-(2-phenyl-[1,3]dioxolan-2-yl)-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 30
- 4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethoxy]-phenol,
- 35
- [4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-phenyl-methanone,

- 5-[2-(4-Fluoro-phenoxy-methyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-[2-(2,2-Dimethyl-propyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[2-(4-Fluoro-phenylsulfanylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 10 [4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-yl]-methanol,
- 4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-carbaldehyde
- 5-[2-[(4-Fluoro-phenylamino)-methyl]-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 15 5-(2-Imidazol-1-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 20 5-(2-Dimethylaminomethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(6,7-Dimethoxy-2-morpholin-4-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 25 5-(2-Biphenyl-4-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid,
- 30 4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-3-methyl-benzofuran-2-carboxylic acid ethyl ester,
- 35 5-[6,7-Dimethoxy-2-((4-chlorobenzyl)-benzofuran-4-ylmethyl)pyrimidine-2,4-diamine,

- 5-[6,7-Dimethoxy-2-(1-naphthylmethyl)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(2-propenyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5 5-(6,7-Dimethoxy-2-trifluoromethylbenzofuran-4-ylmethyl)pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(2,2-dimethylpropan-1-one)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 10 5-[6,7-Dimethoxy-2-(cyclopropylcarbonyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(4-methoxyphenyl-methanone)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 15 5-[6,7-Dimethoxy-2-(4-chlorophenyl-methanone)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(4-fluorophenyl-methanone)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 20 5-[6,7-Dimethoxy-2-(1-naphthyl-methanone)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 25 5-[6,7-Dimethoxy-2-(2,2-dimethyl-1-hydroxypropyl)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(cyclopropylmethanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 30 5-[6,7-Dimethoxy-2-(phenylmethanol)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(4-methoxyphenylmethanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 35

- 5-[6,7-Dimethoxy-2-((4-chlorophenyl)methanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-((4-fluorophenyl)methanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(1-naphthylmethanol)-benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 10 5-[2-(3,4-Dihydro-2H-quinolin-1-ylmethyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(tetrazol-5-ylmethyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 15 5-[6,7-Dimethoxy-2-(indol-1-ylmethyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 5-[6,7-Dimethoxy-2-(imidazol-1-ylcarbonyl)benzofuran-4-ylmethyl]pyrimidine-2,4-diamine,
- 20 5-(2-Furan-2-ylmethyl-6,7-dimethoxy-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(6,7-Dimethoxy-2-thiophen-2-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 25 5-[6,7-Dimethoxy-2-(1-methyl-1H-pyrrol-2-ylmethyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,
- 5-(6,7-Dimethoxy-2-pyridin-2-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 30 5-(6,7-Dimethoxy-2-pyridin-3-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 5-(6,7-Dimethoxy-2-pyridin-4-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,
- 35 4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-phenol,

N-{4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-phenyl}-acetamide,

5-[2-(4-Amino-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,

5

5-[6,7-Dimethoxy-2-(4-nitro-benzyl)-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,

N-{4-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-2-methoxy-phenyl}-acetamide,

10

5-[2-(4-Amino-3-methoxy-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,

N-{5-[4-(2,4-Diamino-pyrimidin-5-ylmethyl)-6,7-dimethoxy-benzofuran-2-ylmethyl]-2-methoxy-phenyl}-acetamide,

15

5-[2-(3-Amino-4-methoxy-benzyl)-6,7-dimethoxy-benzofuran-4-ylmethyl]-pyrimidine-2,4-diamine,

20 5-(6,7-Dimethoxy-2-thiophen-3-ylmethyl-benzofuran-4-ylmethyl)-pyrimidine-2,4-diamine,

and pharmaceutically acceptable salts and N-oxides thereof.

25 7. Pharmaceutical compositions for the treatment of infections containing a compound of any one of claims 1 to 6 and usual carrier materials and adjuvants.

8. Pharmaceutical compositions for the treatment of infections caused by Gram positive and Gram negative pathogens, containing a compound of any one of claims
30 1 to 6 and usual carrier materials and adjuvants.

9. The compounds of any one of the claims 1 to 6 for use as medicaments for the treatment of infections.

35 10. The compounds of any one of the claims 1 to 6 for use as medicaments for the treatment of infections caused by Gram positive and Gram negative pathogens.

11. The use of one or more compounds of any one of claims 1 to 6 as active ingredients for the production of pharmaceutical compositions for the treatment of infections.

5

12. The use of one or more compounds of any one of claims 1 to 6 as active ingredients for the production of pharmaceutical compositions for the treatment of infections caused by Gram positive and Gram negative pathogens.

10 13. A process for the manufacture of pharmaceutical compositions for the treatment of infections containing one or more compounds as claimed in any one of claims 1 to 6 as active ingredients which process comprises mixing one or more active ingredient with pharmaceutically acceptable excipients in a manner known per se.

15 14. A process for the manufacture of pharmaceutical compositions for the treatment of infections caused by Gram positive and Gram negative pathogens containing one or more compounds as claimed in any one of claims 1 to 6 as active ingredients which process comprises mixing one or more active ingredient with pharmaceutically acceptable excipients in a manner known per se.

20

15. The invention as hereinbefore described.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 01/08426

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D405/06 C07D405/14 A61K31/505 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 096 214 A (WELLCOME) 21 December 1983 (1983-12-21) cited in the application claims; examples 25-27	1-15
Y	GB 875 562 A (WELLCOME) 23 August 1961 (1961-08-23) cited in the application page 2, line 13 - line 20; claims; examples	1-15
Y	US 5 773 446 A (MASCIADRI) 30 June 1998 (1998-06-30) column 1, line 20 - line 30; claims; examples	1-15
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

12 December 2001

Date of mailing of the international search report

19/12/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/EP 01/08426

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 438 267 A (DALUGE ET. AL.) 20 March 1984 (1984-03-20) column 6, line 38 - line 46; claims; examples -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/EP 01/08426

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0096214	A	21-12-1983	AU 569468 B2	04-02-1988
			AU 1410583 A	03-11-1983
			DE 3382172 D1	04-04-1991
			DK 190983 A	02-11-1983
			EP 0096214 A1	21-12-1983
			FI 831480 A ,B,	02-11-1983
			FI 854491 A	14-11-1985
			HU 195207 B	28-04-1988
			JP 58208277 A	03-12-1983
			KR 9101006 B1	19-02-1991
			US 4590271 A	20-05-1986
			ZA 8303067 A	24-12-1984
			US 4587341 A	06-05-1986
GB 875562	A	23-08-1961	CH 366835 A	31-01-1963
US 5773446	A	30-06-1998	AU 708578 B2	05-08-1999
			AU 7696396 A	27-06-1997
			BR 9611871 A	17-02-1999
			CA 2238521 A1	12-06-1997
			CN 1203600 A	30-12-1998
			WO 9720839 A1	12-06-1997
			EP 1149834 A1	31-10-2001
			EP 0866791 A1	30-09-1998
			JP 2000501399 T	08-02-2000
			TR 9801014 T2	21-09-1998
US 4438267	A	20-03-1984	AT 384023 B	25-09-1987
			AT 483781 A	15-02-1987
			AU 549449 B2	30-01-1986
			AU 7733481 A	20-05-1982
			CA 1186310 A1	30-04-1985
			CS 8108257 A2	12-07-1990
			CY 1419 A	22-04-1988
			DD 201896 A5	17-08-1983
			DE 3173752 D1	20-03-1986
			DK 496681 A	12-05-1982
			EG 16107 A	30-10-1987
			EP 0051879 A2	19-05-1982
			ES 506969 D0	01-07-1983
			ES 8307759 A1	01-11-1983
			ES 521201 D0	01-06-1984
			ES 8405377 A1	16-09-1984
			ES 521202 D0	01-06-1984
			ES 8405378 A1	16-09-1984
			ES 521203 D0	01-06-1984
			ES 8405379 A1	16-09-1984
			ES 521204 D0	01-06-1984
			ES 8405380 A1	16-09-1984
			FI 813546 A ,B,	12-05-1982
			GB 2087881 A ,B	03-06-1982
			GR 81317 A1	11-12-1984
			HK 23388 A	08-04-1988
			HU 188762 B	28-05-1986
			IE 52128 B1	24-06-1987
			IL 64256 A	31-01-1988
			IL 75634 A	31-01-1988
			IL 75635 A	31-01-1988

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/08426

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4438267	A	IT 1210589 B	14-09-1989
		JP 1783158 C	31-08-1993
		JP 4074351 B	26-11-1992
		JP 57114581 A	16-07-1982
		KE 3650 A	01-08-1986
		KR 8801736 B1	10-09-1988
		MC 1418 A	18-10-1982
		MY 67886 A	31-12-1986
		NO 813804 A ,B,	12-05-1982
		NZ 198932 A	31-05-1985
		PH 22654 A	28-10-1988
		PH 25053 A	28-01-1991
		PL 233754 A1	01-08-1983
		PL 239278 A1	01-08-1983
		PL 239279 A1	01-08-1983
		PL 239280 A1	01-08-1983
		PT 73960 A ,B	01-12-1981
		SG 62686 G	27-03-1987
		SU 1424732 A3	15-09-1988